

## **Hair Reduction Treatment**

## Informed Consent Form

I,	, authorize	, and /or a
designated practitioner of Elevate M area(s) of my body:	edical Spa to perform hair reductio	on with BBL on the following
Laser and/or BBL (BroadBand Ligh destroying them while leaving the sudiminish or remove unwanted hair.	, , ,	

## **Review of facts about light therapy**

- Hair reduction requires more than one treatment and does not remove all hair. Generally, hair reduction is decreased in the amount of hair growth by 60-90% on average. Results depend on color and location of hair follicles. A general range of 4-12 treatments spaced 4-8 weeks apart is possible for maximum results.
- Light from a laser can be harmful to eyes and wearing special safety eyewear is necessary at all times during the procedures.
- Light from BBL is an intense burst of light and even though the special safety eyewear is in place, you all sense light emanating from the treatment area.
- The sensation of light may be uncomfortable in certain areas and feel like pin pricks or bursts of heat. The use of topical anesthetics is at the discretion of the practitioner as there are known severe allergic reactions to ingredients in topical anesthetics. Patients with known allergies to anesthetics will list them here:

## Common side effects and risks

- Erythema (redness) may occur in the area of treatment. This may last several hours. Edema (swelling) around the hair follicles is called peri-follicular edema and is a sign that the hair follicle has been affected. Urticaria (itching) or hive-like appearance is also associated with the thermal light affecting the surrounding skin. These symptoms usually subside in a few hours. A cool compress placed on the area provides comfort. The treated area should be cared for delicately for at least 12 hours. Limited activity may be advised, as well as no hot tub, steam, sauna, or shower use.
- A blister can form up to 48 hours after treatment. An antibiotic cream or ointment can be used. Other short term effects include bruising, superficial crusting, and discomfort.
- Hyperpigmentation (browning) and hypopigmentation (lightening) have been noted. These conditions usually resolve within 2-6 months. Permanent color change is a rare risk. Vigilant care must be taken



- to avoid sun exposure (tanning beds included) before and after the treatments to reduce the risk of color change. Sunscreen and / or sun block should be applied when sun exposure is necessary.
- Infection is not usual after treatment; however, herpes simplex virus infections around the mouth can occur following treatments. This applies to both individuals with a past history of the virus or individuals with no known history. Should any kind of infection occur, your clinician must be notified to prescribe appropriate medical care.
- Allergic reactions resulting from treatment are uncommon. Some persons may have a hive-like
  appearance in the treated area as discussed above. Some persons have localized reactions to cosmetics
  or topical preparations. Systemic reactions are rare.

I have read and understand the first and second section: initial
--

- Alternative methods of hair reduction are: shaving, waxing, electrolysis, chemical epilation, and threading. I choose to try hair reduction therapy by laser and/or BBL light therapy.
- I understand that compliance with pre and post care instructions is crucial for success of hair reduction therapy and to prevent unnecessary side effects or complications.
- I understand that the hair reduction therapy involves payment and the fee structure has been explained to me.

I have read and understand all information presented to me before signing this consent form. I have been given an opportunity to have all of my questions answered to my satisfaction. I understand the procedure and accept the risks. I agree to the terms of this agreement.

Patient's Name (Printed): _	 	 
Signature:		 
Date:		
Witness:		